



Vol 5 / Issue 2 / April-June 2015

# International Journal of Critical Illness & Injury Science

*Connecting Cells, Clinics and Communities across Continents*

Official Publication of International Network of  
Critical Illness and Injury Trial Experts

[www.ijciis.org](http://www.ijciis.org)

## Sickle cell trait at high altitude

Dear Editor,

Sickle cell anemia (SCA) is one of the common hemoglobinopathies in the world. It can affect any part of the body and one of the most common and an early organ to be affected in SCA is the spleen. Massive splenic infarction which is arbitrarily defined as infarction involving more than 50% of the spleen size is a rare and unique complication of SCA.<sup>[1]</sup> In sickle cell trait, large splenic infarcts in the normal sized spleen were reported following exposure to hypoxia during high altitude flights in an unpressurized air planes or during mountain climbing.<sup>[2,3]</sup>

Herein, we present a case of sickle cell trait associated with splenic infarction after mountain climbing. An 18-year-old male presented to our emergency department because of severe left upper quadrant pain and fever after mountain climbing. There was no history of similar pain before. In Physical examination, conjunctiva was pale and oral temperature was 38.1°C. Para clinic tests revealed thrombocytopenia, high serum lactate dehydrogenase (LDH) level and mild elevation in aspartate aminotransferase (AST) and bilirubin. A contrast-enhanced computed tomography (CT) scans of the abdomen and pelvis were consistent with




**Figure 1:** A contrast enhanced computed tomography (CT) scan of the abdomen and pelvis was consistent with splenic infarction

splenic infarction [Figure 1]. According to history of pain after mountain climbing and splenic infarction, SCA was the most probable diagnosis. Hemoglobin electrophoresis and gene analysis were consistent with sickle cell trait [Figure 2]. Treatment with nasal oxygen and hydration was started. Severe pain was controlled with opioid analgesics and was relieved gradually without any complication such as splenic bleeding or rupture. Splenic infarcts had resolved gradually, without any specific treatment or surgical intervention.

In conclusion, physicians should consider splenic infarction in patients who develop suspicious symptoms after exposure to a high altitude environment. These patients could respond well to supportive management, and splenectomy would be avoided.

Shahram Habibzadeh, Nasrollah Maleki<sup>1</sup>

Departments of Infectious Disease and Tropical Medicine, and <sup>1</sup>Internal Medicine, Imam Khomeini Hospital, Ardabil University of Medical Science, Ardabil, Iran



**Figure 2:** Hemoglobin electrophoresis was consistent with sickle cell trait

Address for correspondence:

Dr. Nasrollah Maleki,

Department of Internal Medicine, Imam Khomeini Hospital,

Ardabil University of Medical Sciences, Ardabil, Iran.

E-mail: malekinasrollah@yahoo.com

## REFERENCES

1. Al Jama AH, Al Salem AH, Al Dabbous IA. Massive splenic infarction in Saudi patients with sickle cell anemia: A unique manifestation. *Am J Hematol* 2002;69:205-9.
2. Stock AE. Splenic infarction associated with high altitude flying and sickle cell trait. *Ann Intern Med* 1956;44:554-6.
3. O'Brien RT, Pearson HA, Godley JA, Spencer RP. Splenic infarction and sickle-(cell) trait. *N Engl J Med* 1972;287:720.

## Access this article online

Website: [www.ijciis.org](http://www.ijciis.org)

DOI: 10.4103/2229-5151.158428

Quick Response Code:

